

7N-19-CR  
176496

1N  
19  
—

**FINAL REPORT**

p-32

**Analysis of Satellite Data for Sensor Improvement**

**NASA Grant NGR 14-001-008**

**01 October 1962 to 30 June 1990**

(NASA-CR-193329) ANALYSIS OF  
SATELLITE DATA FOR SENSOR  
IMPROVEMENT Final Report, 1 Oct.  
1962 to 30 Jun. 1990 (Chicago  
Univ.) 32 p

N93-72595

Unclass

29/19 0176496

**Tetsuya Theodore Fujita**

**The University of Chicago**

**5734 S. Ellis Avenue  
Chicago, Illinois 60637**

## GENERAL STATEMENT

This grant began shortly after the successful launch of TIROS 1 on April 1, 1960. Fujita was called into NASA to solve their gridding and navigation problems. For two to three years after this grant started, Fujita and his associates solved the gridding method, making use of the celestial coordinates of the spin-axis point.

After the NIMBUS satellite was launched, Fujita and Bandeen collaborated on identifying successive scan lines on the earth, enabling us to locate precisely the scan spots on the earth. Meanwhile, enhancement of radiation data was suggested.

When the gridding accuracy improved, meteorological applications of satellite were explored. The hurricane structure and effects of seeding experiments were studied in cooperation with NHRP (National Hurricane Research Project).

In cooperation with Suomi and Parents of the University of Wisconsin, the "First Color Movie of the Planet Earth" was produced by using a sequence of color pictures of the earth on November 18, 1967 at 30-min intervals.

Since that time, Fujita developed and improved his technique of making movies of all types of nephosystems such as thunderstorms, jetstream cirrus clouds, etc. A list of ten (10) significant movies produced under this grant is shown in p11 and p12.

1974 was the year of the Super-outbreak tornadoes and the year of "Hearings before the Subcommittee on Space Science and Applications." After the hearing, a large-scale investigation of thunderstorm tops during the tornado-bearing stages was undertaken. Among others, it has been confirmed that cloud tops either subside or collapse prior to tornado development.

When the warm wake atop severe thunderstorms were studied extensively by Fujita, resulting in his hypothesis that the wake is caused by the stratospheric cirrus located above the anvil surface. NASA, on the other hand, maintained their view that the emissivity of cloud tops may as well cause the warm wake. This debate could go on for some years to come.

Extensive research on overshooting tops was also performed by photographing the tops from a Learjet flying above the anvil level. Amazingly enough, the time variation of the dome top height is very fast.

After the discovery of downburst by Fujita, a number of attempts were made for detecting downburst clouds prior to their formation. Unfortunately, small downburst cells failed to show specific signatures in satellite pictures.

During the past years, Fujita began thinking about the climatic change of severe storm activities, say, during the 30 to 60 years when rawinsonde data were archived. Fujita's interest consequently widened into climatological researches.

A W A R D S   R E C E I V E D

- 1979: Distinguished Public Service Medal of NASA to Tetsuya T. Fujita for his significant and imaginative utilization of the measurements from low orbiting and geosynchronous meteorological satellites for nearly two decades, resulting in major advances in our understanding the structure and evolution of severe storms.
- 1985: 25-Years of Weather Satellite Award Medal of the U.S. Department of Commerce to T. Theodore Fujita for creative scientific leadership as an enthusiastic pioneer in the use of satellite imagery to analyze and predict mesoscale weather phenomena and to understand severe thunderstorms, tornadoes, and hurricanes.

Author	Paper Identification Numbers
Arnold, J. E.	5 10
Balogun, E. E.	51
Bandeen, W.	9
Black, P. G.	24 92
Bonner, W. D.	11
Bradbury, D. L.	19
Byers, H. R.	42
Caracena, F.	48 50
Changnon, S. A.	44
Dundus, R.	45
Fletcher, W. L.	2 4
Forbes, G. S.	32 36 40 46 54 75
Fujita, T. T.	1 3 4 5 6 7 8 9 13 16 19 21 23 24 25 26 27 28 30 31 32 33 34 38 39 40 42 43 44 45 47 48 49 50 52 53 56 57 58 59 60 61 62 64 65 66 67 69 70 71 72 76 77 78 80 81 82 84 85 90 91 92 93 94
Grandoso, H.	16
Hjelmfelt, M. J.	44
Larsen, S. H. H.	2 4
Madden, R. A.	18
McGurk, P. D.	55
Mendez, R.	17
Ninomiya, K.	20
Partacz, J. W.	87 93
Pearl, E. W.	29 33 37 41
Pedersen, F.	3
Peterson, E. C.	88
Rabbe, A.	8 14
Reap, R. M.	21 13 15
Shenk, W. E.	33 37
Skilman, W.	37
Smith, B. E.	79 87 93
Stiegler, D. J.	58 60 63 82 85
Tecson, J. J.	22 38 47 49
Umenhofer, T. A.	35 40 45 47
Wakimoto, R. M.	54 58 60 65 68 71 72 73 74 75 77 78 83 86
Waranauskas, B.R.	79 89
Wilson, J. W.	57

P A P E R    B Y    S U B J E C T    M A T T E R S

Subjects	Paper Identification Numbers												
<u>A. THUNDERSTORM</u>	<u>16</u>	19	28	34	42	51	52	55	56	57	67	68	73
	80	79	94										
<u>B. HURRICANE</u>	<u>5</u>	15	24	60	82								
<u>C. TORNADO</u>	<u>20</u>	23	26	27	30	31	32	36	37	44	46	47	53
	54	62	66	71	74	75	85	87	91	94			
<u>D. DOWNBURST</u>	<u>40</u>	43	48	50	52	54	59	61	64	65	66	77	80
	84	86	89	90	92								
<u>E. MESOSCALE FLOW</u>	<u>81</u>	83	88	93									
<u>F. CLOUD MOTION</u>	<u>33</u>	39	48										
<u>G. PRECISE GRIDDING</u>	<u>1</u>	6	7	10	11	18							
<u>H. STEREO</u>	<u>69</u>	70	76										
<u>I. EARTH SCANNING</u>	<u>9</u>	10	17	21									
<u>J. GROUND TRUTH</u>	<u>12</u>	13	49	58	72								
<u>K. STORM TOP</u>	<u>16</u>	25	29	30	35	38	41	51	55	63			
<u>L. RAINSTORM</u>	<u>45</u>												
<u>M. RADIATION</u>	<u>1</u>	2	3	4	5	6	7	8	14	22			

# LIST OF PUBLICATIONS

- 1 Fujita, T., 1963: Outline of a theory and examples for precise analysis of satellite radiation data. MP Res. Paper 15, University of Chicago, 35 pp.
- 2 Larsen, S.H.H., T. Fujita, and W.L. Fletcher, 1963: Evaluation of limb darkening from TIROS III radiation data. MP Res. Paper 18, University of Chicago, 19 pp.
- 3 Pedersen, F., and T. Fujita, 1963: Synoptic interpretation of TIROS III measurements of infrared radiation. MP Res. Paper 19, University of Chicago, 39 pp.
- 4 Larsen, S.H.H., T. Fujita, and W.L. Fletcher, 1963: TIROS III measurements of terrestrial radiation and reflected and scattered solar radiation. MP Res. Paper 20, University of Chicago, 11 pp.
- 5 Fujita, T., and J. Arnold, 1963: The decaying stage of Hurricane Ana of July 1961 as portrayed by TIROS cloud photographs and infrared radiation from the top of the storm. MP Res. Paper 28, University of Chicago, 13 pp.
- 6 Fujita, T., 1964: A technique for precise analysis of satellite data: Volume II - Radiation analysis. SMRP Res. Paper 29, University of Chicago, 20 pp.
- 7 Fujita, T., 1964: The scanning printer and its application to detailed analysis of satellite radiation data. SMRP Res. Paper 34, University of Chicago, 29 pp.
- 8 Rabbe, A., and T. Fujita, 1964: Synoptic study of cold air outbreak over the Mediterranean using satellite photographs and radiation data. SMRP Res. Paper 35, University of Chicago, 33 pp.
- 9 Fujita, T., and W. Bandeen, 1965: Resolution of the NIMBUS high resolution infrared radiometer. SMRP Res. Paper 40, University of Chicago, 11 pp.
- 10 Arnold, J.E., 1965: A simplified grid technique for determining scan lines generated by the TIROS scanning radiometer. SMRP Res. Paper 32, University of Chicago, 18 pp.
- 11 Bonner, W.D., 1965: A program for computer gridding of satellite photographs for mesoscale research. SMRP Res. Paper 43, University of Chicago, 31 pp.
- 12 Reap, R.M. 1965: Comparison of grassland surface temperatures measured by TIROS VII and airborne radiometers under clear sky and cirriform cloud conditions. SMRP Res. Paper 44, University of Chicago, 35 pp.
- 13 Reap, R.M., and T. Fujita, 1965: Death Valley temperature analysis utilizing NIMBUS I infrared data and ground-based measurements. SMRP Res. Paper 45, University of Chicago, 19 pp.
- 14 Rabbe, A., 1966: Angular dependence of albedo from stratiform clouds as measured by TIROS IV scanning radiometers. SMRP Res. Paper 58, University of Chicago, 19 pp.
- 15 Reap, R.M., 1966: Preliminary investigation of peripheral subsidence associated with hurricane outflow. SMRP Res. Paper 63, University of Chicago, 19 pp.

- 16 Fujita, T., and H. Grandoso, 1967: A proposed method of estimating cloud-top temperature, cloud covers, emissivity, and cloudiness from short- and long-wave radiation data obtained by medium resolution scanning radiometers. SMRP Res. Paper 48, University of Chicago, 48 pp.
- 17 Mendez, R., 1968: Angular dependence of reflected solar radiation from Sahara measured by TIROS VII in a torquing maneuver. SMRP Res. Paper 73, University of Chicago, 17 pp.
- 18 Madden, R.A., 1969: Yaw corrections for accurate gridding of NIMBUS HRIR data. SMRP Res. Paper 77, University of Chicago, 26 pp.
- 19 Fujita, T.T., and D.L. Bradbury, 1969: Determination of mass outflow from a thunderstorm complex using ATS III pictures. SMRP Res. Paper 79, University of Chicago, 15 pp.
- 20 Ninomiya, K., 1969: Dynamical analysis of outflow from tornado-producing thunderstorms as revealed by ATS III pictures. SMRP Res. Paper 81, University of Chicago, 43 pp.
- 21 Fujita, T.T., 1970: Basic problems on cloud identification related to the design of SMS-GOES spin scan radiometers. SMRP Res. Paper 84, University of Chicago, 33 pp.
- 22 Tecson, J.J., 1970: Patterns of equivalent blackbody temperature and reflectance of model clouds computed by changing radiometer's field of view. SMRP Res. Paper 87, University of Chicago, 27 pp.
- 23 Fujita, T.T., 1970: Lubbock tornadoes of 11 May 1970. SMRP Res. Paper 88, University of Chicago, 23 pp.
- 24 Fujita, T.T., and P.G. Black, 1970: In- and outflow field of Hurricane Debbie as revealed by echo and cloud velocities from airborne radar and ATS-III pictures. Preprints, 14th Radar Meteorology Conf., Tucson, 353-358.
- 25 Fujita, T.T., 1972: Tornado occurrences related to overshooting cloud-top heights as determined from ATS pictures. SMRP Res. Paper 97, University of Chicago, 32 pp.
- 26 Fujita, T.T., 1972: F-scale classification of 1971 tornadoes. SMRP Res. Paper 100, University of Chicago, 16 pp.
- 27 Fujita, T.T., 1973: Proposed mechanism of tornado formation from rotating thunderstorm. Preprints, 8th Conf. on Severe Local Storms, Boston, 191-196.
- 28 Fujita, T.T., 1973: Thermal and dynamical features of a thunderstorm with a tilted axis of rotation. SMRP Res. Paper 118, University of Chicago, 15 pp.
- 29 Pearl, E.W., 1974: Characteristics of anvil-top associated with the Poplar Bluff tornado of May 7, 1973. SMRP Res. Paper 119, University of Chicago, 12 pp.
- 30 Fujita, T.T., 1974: Overshooting thunderheads observed from ATS and Learjet. SMRP Res. Paper 117, University of Chicago, 29 pp.
- 31 Fujita, T.T., 1974: Jumbo tornado outbreak of 3 April 1974. Weatherwise, 27, 116-126.

- 32 Fujita, T.T., and G.S. Forbes, 1974: Superoutbreak tornadoes of April 3, 1974 as seen in ATS pictures. Preprints, 6th Conf. on Aerospace and Aeronautical Meteorology, El Paso, 165-172.
- 33 Fujita, T.T., E.W. Pearl, and W.E. Shenk, 1975: Satellite tracked cumulus velocities, J. Appl. Meteor., 14, 407-413.
- 34 Fujita, T.T., 1975: Detection of severe storms by satellite. Preprints, 3rd Symposium on Meteorological Instruments and Instrumentation, Washington, D.C., 5-9.
- 35 Umenhofer, T.A., 1975: Overshooting top behavior of three tornado-producing thunderstorms. Preprints, 9th Conf. on Severe Local Storms, Norman, 96-99.
- 36 Forbes, G.S., 1975: Relationship between tornadoes and hook echoes on April 3, 1974. Preprints, 9th Conf. on Severe Local Storms, Norman, 280-285.
- 37 Pearl, E.W., W.E. Shenk, and W. Skillman, 1975: Cloud-top parameters - A hail indicator. Preprints, 9th Conf. on Severe Local Storms, Norman, 464-467.
- 38 Tecson, J.J., and T.T. Fujita, 1975: Cloud-motion vectors over the GATE area computed by McIDAS and METRACOM methods. SMRP Res. Paper 136, University of Chicago, 23 pp.
- 39 Fujita, T.T., 1976: Spearhead echo and downburst near the approach end of a John F. Kennedy Airport runway, New York City. SMRP Res. Paper 137, University of Chicago, 51 pp.
- 40 Fujita, T.T., G.S. Forbes, and T.A. Umenhofer, 1976: Close-up view of 20 March 1976 tornadoes: Sinking cloud tops and suction vortices. Weatherwise, 29, 116-131.
- 41 Pearl, E.W., 1976: Feeder clouds associated with tornado-producing mesocyclone of March 26, 1976. SMRP Res. Paper 143, University of Chicago, 18 pp.
- 42 Fujita, T.T., and H.R. Byers, 1977: Spearhead echo and downburst in the crash of an airliner. Mon. Wea. Rev., 105, 129-146.
- 43 Fujita, T.T., 1977: Anticyclonic tornadoes. Weatherwise, 30, 51-64.
- 44 Fujita, T.T., M.J. Hjelmfelt, and S.A. Changnon, 1977: Mesoanalysis of record Chicago rainstorm using radar, satellite, and raingauge data. Preprints, 10th Conf. on Severe Local Storms, Omaha, 65-72.
- 45 Umenhofer, T.A., T.T. Fujita, and R. Dundas, 1977: Great Bend tornadoes of August 30, 1974. Preprints, 10th Conf. on Severe Local Storms, Omaha, 457-462.
- 46 Forbes, G.S., 1977: Thunderstorm-scale variations of echoes associated with left-turn tornado families. Preprints, 10th Conf. on Severe Local Storms, Omaha, 497-504.
- 47 Tecson, J.J., T.A. Umenhofer, and T.T. Fujita, 1977: Thunderstorm-associated cloud motions as computed from 5-minute SMS pictures. Preprints, 10th Conf. on Severe Local Storms, Omaha, 22-29.



- 48 Fujita, T.T., and F. Caracena, 1977: Common denominator of three weather-related aircraft accidents. Preprints, 10th Conf. on Severe Local Storms, Omaha, 135-142.
- 49 Fujita, T.T., and J.J. Tecson, 1977: Mesoscale wake clouds in Skylab photographs. Skylab Explores the Earth, NASA, 463-477.
- 50 Fujita, T.T., and F. Caracena, 1977: An analysis of three weather-related aircraft accidents. Bull. Amer. Meteor. Soc., 58, 1164-1181.
- 51 Balogun, E.E., 1977: A kinematic investigation of the influence of anvil airflow over convective nephsystems on high-level horizontal flow. SMRP Res. Paper 160, University of Chicago, 18 pp.
- 52 Fujita, T.T., 1978: Manual of downburst identification for Project NIMROD. SMRP Res. Paper 156, University of Chicago, 104 pp. [NTIS PB-286048].
- 53 Fujita, T.T., and J.R. MacDonald, 1978: Tornado damage at the Grand Gulf Mississippi nuclear power plant site: Aerial and ground surveys. NUREG/CR-0383, 61 pp.
- 54 Forbes, G.S., and R.M. Wakimoto, 1978: Downbursts and tornadoes of 6 August 1977 near Springfield, Illinois. SMRP Res. Paper 161, University of Chicago, 50 pp. [NTIS PB80-159791].
- 55 McGurk, P.D., 1979: Rawinsonde detection of stratospheric moisture in the vicinity of thunderstorm tops. SMRP Res. Paper 176, University of Chicago, 18 pp.
- 56 Fujita, T.T., 1979: Objectives, operation, and results of Project NIMROD. Preprints, 11th Conf. on Severe Local Storms, Kansas City, 259-266.
- 57 Wilson, J.W., and T.T. Fujita, 1979: Vertical cross section through a rotating thunderstorm by Doppler radar. Preprints, 11th Conf. on Severe Local Storms, Kansas City, 447-452.
- 58 Stiegler, D.J., T.T. Fujita, and R.M. Wakimoto, 1979: The effect of side lobes upon measured Doppler velocities in weak echo regions. Preprints, 11th Conf. on Severe Local Storms, Kansas City, 509-514.
- 59 Fujita, T.T., 1980: Downbursts and microbursts--An aviation hazard. Preprints, 19th Conf. on Radar Meteorology, Miami, 94-101.
- 60 Fujita, T.T., R.M. Wakimoto, and D.J. Stiegler, 1980: Mesoscale damage patterns of Hurricane Frederic in relation to enhanced SMS imagery. Preprints, 19th Conf. on Radar Meteorology, Miami, 48-55.
- 61 Fujita, T.T., 1980: Meso-scale structure of downbursts depicted by Doppler radar. (Supplement to Downburst and microburst--An aviation hazard). SMRP Res. Paper 181, University of Chicago, 19 pp.
- 62 Fujita, T.T., 1980: Diabatic analysis: A new method for predicting tornado outbreaks. SMRP Res. Paper 183, University of Chicago, 15 pp.
- 63 Stiegler, D.J., 1980: Study of a splitting anvil over Iowa on May 15, 1977, using 3-minute SMS and Lear jet pictures. SMRP Res. Paper 168, University of Chicago, 26 pp.
- 64 Fujita, T.T., 1981: Microburst as an aviation wind shear hazard. Preprints, AIAA 19th Aerospace Meeting, St. Louis, 1-9.
- 65 Fujita, T.T., and R.M. Wakimoto, 1981: Five scales of airflow associated with a series of downbursts on 16 July 1980. Mon. Wea. Rev., 19, 1438-1456.

- 66 Fujita, T.T., 1981: Tornadoes and downbursts in the context of generalized planetary scales. J. Atmos. Sci., 38, 1511-1534.
- 67 Fujita, T.T., 1981: Mesoscale aspects of convective storms. SMRP Res. Paper 191, University of Chicago, 8 pp.
- 68 Wakimoto, R.M., 1982: The life cycle of thunderstorm gust fronts as viewed with Doppler radar and rawinsonde data. Preprints, 12th Conf. on Severe Local Storms, San Antonio, 409-412.
- 69 Fujita, T.T., 1982: Principle of stereoscopic height computations and their applications to stratospheric cirrus over severe thunderstorms. J. Meteor. Soc. Japan, 60, 355-368.
- 70 Fujita, T.T., 1982: Infrared, stereo-height, cloud-motion, and radar-echo analysis of Sesame-Day thunderstorms. Preprints, 12th Conf. on Severe Local Storms, San Antonio, 213-216.
- 71 Fujita, T.T., and R.M. Wakimoto, 1982: Anticyclonic tornadoes in 1980 and 1981. Preprints, 12th Conf. on Severe Local Storms, San Antonio, 401-404.
- 72 Fujita, T.T., and R.M. Wakimoto, 1982: Effect of miso- and mesoscale obstructions on PAM winds obtained during Project NIMROD. J. Appl. Meteor., 21, 840-858.
- 73 Wakimoto, R.M., 1982: The life cycle of thunderstorm gust fronts as viewed with Doppler radar and rawinsonde data. Mon. Wea. Rev., 110, 1060-1082.
- 74 Wakimoto, R.M., 1983: The West Bend, Wisconsin storm of 4 April 1981: A problem in operational meteorology. J. Climate and Appl. Meteor., 22, 181-189.
- 75 Forbes, G.S., and R.M. Wakimoto, 1983: A concentrated outbreak of tornadoes, downbursts and microbursts, and implications regarding vortex classification. Mon. Wea. Rev., 111, 220-235.
- 76 Fujita, T.T., and J.C. Dodge, 1983: Applications of stereoscopic height computations from dual geosynchronous satellite data/Joint NASA-Japan stereo project. Ad. Space Res., 2, 153-160.
- 77 Fujita, T.T., and R.M. Wakimoto, 1983: Microbursts in JAWS depicted by Doppler radars, PAM, and aerial photographs. Preprints, 21st Conf. on Radar Meteorology, Edmonton, 638-645.
- 78 Fujita, T.T., and R.M. Wakimoto, 1983: JAWS microbursts revealed by triple-Doppler radar, aircraft, and PAM data. Preprints, 13th Conf. on Severe Local Storms, Tulsa, 97-100.
- 79 Smith, B.E., and B.R. Waranauskas, 1983: Analysis of wet microbursts by dual-Doppler and ground photography. Preprints, 13th Conf. on Severe Local Storms, Tulsa, 51-54.
- 80 Fujita, T.T., 1983: Andrews AFB microburst. SMRP Res. Paper 205, University of Chicago, 38 pp.
- 81 Fujita, T.T., 1984: Review of the history of mesoscale meteorology and forecasting. SMRP Res. Paper 208, University of Chicago, 42 pp.

- 82 Stiegler, D.J., and T.T. Fujita, 1984: Mesoscale structure of Hurricane Alicia. Hurricane Alicia: One Year Later, Ashan Kareem, Ed., Amer. Soc. Civ. Engrs., 21-39.
- 83 Wakimoto, R.M., 1984: Clear air mesocyclone during the JAWS Project. Preprints, 22nd Conf. on Radar Meteorology, Zurich, 476-481.
- 84 Fujita, T.T., 1985: The Downburst: Microburst and macroburst. SMRP Res. Paper 210, University of Chicago, 122 pp. Hardbound color book. [NTIS PB85-148880, Library of Congress No. 85-50115].
- 85 Fujita, T.T., and D.J. Stiegler, 1985: Detailed analysis of the tornado outbreak in the Carolinas using radar, satellite, and aerial survey data. Preprints, 14th Conf. on Severe Local Storms, Indianapolis, 271-274.
- 86 Wakimoto, R.M., 1985: Forecasting dry microburst activity over the High Plains. Mon. Wea. Rev., 113, 1131-1143.
- 87 Smith, B.E., and J.W. Partacz, 1985: Bow-induced tornado at Minneapolis on 26 April 1984. Preprints, 14th Conf. on Severe Local Storms, Indianapolis, 81-84.
- 88 Peterson, E.C., 1985: JAWS microbursts occurring from orographic vortex streets. Preprints, 14th Conf. on Severe Local Storms, Indianapolis, 256-259.
- 89 Waranauskas, B.R., 1985: The rotor microburst: A new explanation for burst swath damage. Preprints, 14th Conf. on Severe Local Storms, Indianapolis, 260-263.
- 90 Fujita, T.T., 1986: DFW microburst. SMRP Res. Paper 217, University of Chicago, 154 pp. Hardbound color book. [NTIS PB86-131638, Library of Congress No. 85-52386].
- 91 Fujita, T.T., 1987: U.S. Tornadoes, Part 1, 70-Year statistics. SMRP Res. Paper 218, University of Chicago, 122 pp. Hardbound color book. [NTIS PB87-127742, Library of Congress No. 86-51637].
- 92 Fujita, T.T., and P.G. Black, 1988: Monrovia microburst of 20 July 1986: A study of "SST." Preprints, 15th Conf. on Severe Local Storms, Baltimore, 380-383.
- 93 Smith, B.E., T.T. Fujita, and J.W. Partacz, 1988: A method of correcting PAM winds based on obstructions and measured windspeed. Preprints, 15th Conf. on Severe Local Storms, Baltimore, 339-342.
- 94 Fujita, T.T., 1989: The Teton-Yellowstone tornado of 21 July 1987. Mon. Wea. Rev., 117, 1913-1940.

LIST OF SATELLITE MOVIES

When Applications and Technology Satellite (ATS) III was tested in 1967 and 1968, Fujita developed various methods of examining sequences of imagery in the motion picture form. The first attempt was made by using a series of color pictures of the earth taken on November 18, 1967. Both Professors Suomi and Parents contributed significantly to this effort. A list of key movies filmed at the University of Chicago under this grant is presented with a brief explanation of each movie.

**A. Color Movie of the Planet Earth (400 ft)**

This is the "First Color Movie of the Planet Earth" filmed from the first and only sequence of ATS III color pictures. It was taken every 30 min on November 18, 1967. The film was made by Fujita in cooperation with Professor Verner Suomi at the University of Wisconsin.

**B. Tornado Situations on April 19, 1968 (400 ft)**

During the Spring season in 1968, NASA conducted the Tornado Watch Experiment. This movie was made by adding red dot at the location of each tornado on the satellite picture at the tornado time. The film was made by Fujita with Suomi and Oliver.

**C. Tornado Situations on April 23, 1968 (400 ft)**

A large number of tornadoes occurred over a large area from Illinois to Indiana. This movies was made by adding red tornado dots to depict tornado activities in relation to satellite-viewed nephsystems. The film was made by Fujita with Suomi and Oliver.

**D. Hurricane Watch Experiment, 1968 (400 ft)**

The overall view of Hurricane Abby was filmed for a period of four (4) days from June 3 to 6, 1968. This film was made as a joint effort of Fujita, Suomi, Oliver, and Gentry.

**E. Close-up views from Hurricane Watch 1968 (400 ft)**

Selected views of small areas inside the hurricane watch imagery in 1968 were used to produce high-resolution movie segments.

JUN 03, 1968	Drifting anvil from Hurricane Abby Sea breeze over the Yucatan Peninsula Arc clouds from thunderstorm areas
JUN 05, 1968	Cloud bands
JUN 22, 1968	Cumulonimbi over Cuba
JUN 23, 1968	Cumulonimbi over Florida

#### **F. Enhanced Imagery of Hurricane Camille, AUG 17, 1969 (200 ft)**

The hurricane imagery was enhanced so that the coldest cloud areas are 100% white and the warmer clouds, graduated grey to black. This movie shows excellent 3-dimensional effects. Meanwhile, the identical enhancement was applied to Hurricane Debby, August 20, 1969 imagery to produce Debby's movie. Finally, the coldest areas were enhanced black in order to produce imagery with black cold top.

#### **G. Hurricane Ginger, SEP 25 - 28, 1971 (180 ft)**

This hurricane was seeded by NHRL (National Hurricane Research Laboratory) aircraft on alternate days of the experiment: September 25 (no seeding), September 26 (seeded), September 27 (no seeding), and September 28 (seeded). The purpose of the film is to depict the difference in the time-lapse view of a seeded hurricane. Red dots are added on the satellite pictures at the time and location of the seeding aircraft. This was a joint effort of Fujita and Gentry.

#### **H. Overshooting Tops of Thunderstorms May 12, 1972 (180 ft)**

Severe thunderstorm clouds were depicted by three scales of motion as viewed at ATS III spin scan camera, and a time-lapse movie camera operated by Fujita on board a research Learjet flying over southern Texas at 47,000 ft pressure altitude. The satellite movie shows the parent thunderstorm and gust fronts radiating out from convective areas.

Detailed motions of the overshooting tops were time-lapsed and re-filmed into 1 sec and 6 sec sequences to show accelerated views of the top of the severe thunderstorm.

#### **I. Tornado Overflight, FEB 23, 1977 (130 ft)**


This film shows the track of a research Learjet on satellite pictures, followed by the sequence of views of a growing overshooting top and a collapsing top, beneath which an F2 tornado was in progress. This film documents that the top of the thunderstorm during its tornado-bearing stage is suppressed. The Learjet was manned by Fujita, Forbes, Umenhofer, and Stiegler of the University of Chicago.

#### **J. Summary of ATS III Cloud Movies (250 ft)**

This movie was made by combining several scenes filmed between 1967 and 1969. These scenes are:

NOV 18, 1967	Color movie of the planet earth
APR 03, 1968	Fishbone cirrus in jet stream
APR 19, 1968	Tornado situations
APR 23, 1968	Tornado situations
MAY 24, 1968	Jet stream cirrus on moving coordinates
JUN 21, 1968	Hurricane outflow
JUN 22, 1968	Hurricane Candy
FEB 23, 1969	Hook-shaped stratus cloud in satellite movie
APR 23, 1969	Source region of subtropical jet cirrus over the Pacific
AUG 17, 1969	Hurricane Camille in regular and enhanced satellite pictures

### Specific Sites of Computing Tornado Statistics




Forty cities selected in computing the long-term variation of tornadoes. Both touchdown counts and path lengths within  $1.0^\circ$  latitude and  $0.75^\circ$  longitude from each city were determined from the University of Chicago Tornado Tape being updated under this NASA Grant, NGR 14-001-008. Included in this final report are nine (9) cities, Chicago, IL; Dallas/Ft Worth TX, Denver, CO, Houston, TX, Huntsville, AL; Kansas City, MO, Oklahoma City, OK; Orlando, FL and Washington D.C.

Diagrams for each city are: Tornado counts in four seasons between 1950 and 1988. Tornado in each year from 1916 to 1988. In order to expand this research into that of "Climatological variation of severe storms, a continuing research proposal was submitted to NASA.

## CHICAGO IL


Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote

Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes




## CHICAGO IL

## PATH LENGTH



## TOUCHDOWN COUNT






# DALLAS/FT WORTH TX


Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote

Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes




## DALLAS/FT. WORTH TX

## PATH LENGTH




## TOUCHDOWN COUNT



## DENVER CO

Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote

Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## DENVER CO


PATH LENGTH

KM PER 10.000 SQ KM



TOUCHDOWN COUNT


PER 10.000 SQ KM



## HOUSTON TX

Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote


Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## HOUSTON TX


PATH LENGTH

KM PER 10.000 SQ KM



TOUCHDOWN COUNT


PER 10.000 SQ KM



# HUNTSVILLE AL

Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote


Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## HUNSTVILLE AL


PATH LENGTH

KM PER 10.000 SQ KM



TOUCHDOWN COUNT

PER 10.000 SQ KM






# KANSAS CITY

Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote


Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## KANSAS CITY MO


PATH LENGTH

KM PER 10.000 SQ KM



TOUCHDOWN COUNT


PER 10.000 SQ KM



# OKLAHOMA CITY OK

Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote


Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## OKLAHOMA CITY OK


PATH LENGTH

KM PER 10.000 SQ KM



TOUCHDOWN COUNT


PER 10.000 SQ KM



## ORLANDO FL

Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote


Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## ORLANDO FL


PATH LENGTH

KM PER 10.000 SQ KM



TOUCHDOWN COUNT


PER 10.000 SQ KM



## WASHINGTON DC


Touchdown counts of tornadoes between 1950 and 1988 by season in each year. Season identifications are W -- Winter, S -- Spring, S -- Summer, and A -- Autumn. Three thickness lines denote

Thin Line F0 and F1 tornadoes  
 Medium Line F2 and F3 tornadoes  
 Heavy Line F4 and F5 tornadoes



## WASHINGTON DC

## PATH LENGTH



## TOUCHDOWN COUNT

